# CHICK DYSENTERY CARE

**(An organic remedy for any intestinal infection)**

**Indication:** Organic remedy for Coccidiosis, diarrhea and blood dysentery for chicken only.

**Benefits:**

- Helps to control of intestinal and caecal coccidiosis in chicken.
- Helps to prevent dehydration and blood loss.
- Helps to cure Coccidiosis, diarrhea and blood dysentery.
- Helps to compact the watery stool.
- Helps to replenish normal intestinal micro flora.
- Helps to digest food and absorbs nutrient.
- Helps to increases appetite & promotes growth.
- Extremely safe and have no side effect.
- No requirement to use of any antibiotics.

**Dosage:**

- ✓ If the intestinal infection are not fully controlled by using of SUBITO PLUS within 12-24 hours then two tea spoon of powder should be mixed with 2-3 litre of mineral water and should be provided for 1000 birds in poultry drinkers in the morning only. Remaining medicine could be used as per direction of a physician.

**Precautions:**

- This medicine should be provided in the morning only.
- Medicine should be used within one month after opening the cap.
- Store in a dry place at a room temperature and keep distant from children.

Organic Research Food Products Ltd.
Pabna, Bangladesh

# চিক ডিসেন্ট্রি কেয়ার

## (অন্ত্রের যে কোনো সংক্রামনের একটি জৈব প্রতিকারক)

**কার্যকারিতা:** শুধুমাত্র মুরগীর কক্সিডিওসিস, ডায়রিয়া ও রক্ত আমাশয় নিরাময়ের একটি জৈব প্রতিকারক।

## উপকারিতা:

- মুরগীর অন্ত্র ও সিকামের কক্সিডিওসিস নিরাময়ে সহায়তা করে।
- পানি শুন্যতা ও রক্তপাত প্রতিরোধ করতে সহায়তা করে।
- কক্সিডিওসিস, ডায়রিয়া ও রক্ত আমাশয় নিরাময়ে সহায়তা করে।
- পানিযুক্ত নরম মলকে শক্ত করতে সহায়তা করে।
- অন্ত্রের মাইক্রো ফ্লোরা পুনরায় পূর্ণ করতে সহায়তা করে।
- খাদ্য পরিপাক ও পুষ্টি শোষন করতে সহায়তা করে।
- ক্ষুধা ও দৈহিক বৃদ্ধিতে সহায়তা করে।
- অত্যন্ত নিরাপদ ও কোন পার্শ্ব প্রতিক্রিয়া নেই।
- কোনো প্রকার এন্টিবায়োটিক ব্যবহারের প্রয়োজন নেই।

## ব্যবহার বিধি:

- ✓ "সুবিতো প্লাস" প্রয়োগের ১২-২৪ ঘন্টার মধ্যে অন্ত্রের সংক্রামন পুরোপুরি নিয়ন্ত্রন না হলে দুই চা চামচ পাউডার ২-৩ লিটার বোতলজাত পানির সাথে মিশিয়ে পানির পাত্রে দিয়ে শুধুমাত্র সকালে ১০০০টি মুরগীর বাচ্চাকে খাওয়াতে হবে। বাকী ঔষধ পরবর্তীতে চিকিৎসকের পরামর্শ অনুযায়ী খাওয়াতে হবে।

Organic Research Food Products Ltd.
Pabna, Bangladesh

# চিক ডিসেন্ট্রি কেয়ার

## (অন্ত্রের যে কোনো সংক্রামনের একটি জৈব প্রতিকারক)

**সতর্কতা:**

- এই ঔষধটি শুধুমাত্র সকালে খাওয়ানো যাবে।
- ছিপি খোলার এক মাসের মধ্যে ঔষধ ব্যবহার করতে হবে।
- ঠান্ডা ও শুষ্ক স্থানে এবং শিশুদের নাগালের বাহিরে রাখুন।