CHICK NAVEL CURE

(An organic Omphalitis or navel-yolk sac infection remedy)

Indication: Organic Omphalitis or navel-yolk sac infection remedy for chicken only.

Benefits:

- Helps to cure any navel deformities of chicks.
- Helps to improve normal droppings.
- Helps to reduce subcutaneous oedema.
- Helps to close the navel opening naturally.
- Helps to increase the body immunity.
- Extremely safe and have no side effect.
- No requirement to use of any antibiotics.

Dosage:

- ✓ Supplied medicine should be used for 1-7 days aged of 1000 birds. Half of supplied powder will be mixed with 100 ml of mineral water and should be provided only one drop orally or should be mixed with 2-3 litre of mineral water in poultry drinkers. Remaining half of medicine could be used next time as per direction of a physician.

Precautions:

- Medicine should be used within one month after opening the cap.
- Store in a dry place at a room temperature and keep distant from children.

Organic Research Food Products Ltd.
Pabna, Bangladesh

চিক ন্যাভেল কিওর

(অম্ফালিটিস বা নাভী-কুসুম থলি সংক্রামনের একটি জৈব প্রতিরোধক)

কার্যকারিতা: শুধুমাত্র মুরগীর নাভী-কুসুম থলি সংক্রামনের একটি জৈব প্রতিরোধক।

উপকারিতা:

- মুরগীর বাচ্চার যে কোনো ধরনের নাভীর বিকৃতি নিরাময়ে সহায়তা করে।
- মলের স্বাভাবিক উন্নয়নে সহায়তা করে।
- ত্বক নিম্নস্থ শোথ কমাতে সহায়তা করে।
- প্রাকৃতিকভাবে নাভী বন্ধ করতে সহায়তা করে।
- শরীরের রোগ প্রতিরোধ ক্ষমতা বৃদ্ধিতে সহায়তা করে।
- অত্যন্ত নিরাপদ ও কোন পার্শ্ব প্রতিক্রিয়া নেই।
- কোনো প্রকার এন্টিবায়োটিক ব্যবহারের প্রয়োজন নেই।

ব্যবহার বিধি:

- ✓ সরবরাহকৃত ঔষধটি ১-৭ দিন বয়সের ১০০০টি মুরগীর বাচ্চাকে খাওয়ানো যাবে। সরবরাহকৃত পাউডারের অর্ধেক ১০০ মিলি বোতলজাত পানির সাথে মিশিয়ে শুধুমাত্র এক ফোটা করে প্রতিটি মুরগীর বাচ্চাকে খাওয়াতে হবে অথবা ২-৩ লিটার বোতলজাত পানির সাথে মিশিয়ে পানির পাত্রে দিয়ে খাওয়াতে হবে। বাকী অর্ধেক ঔষধ পরবর্তীতে চিকিৎসকের পরামর্শ অনুযায়ী খাওয়াতে হবে।

Organic Research Food Products Ltd.
Pabna, Bangladesh

চিক ন্যাভেল কিওর

(অম্ফালিটিস বা নাভী-কুসুম থলি সংক্রামনের একটি জৈব প্রতিরোধক)

সতর্কতা:

- ছিপি খোলার এক মাসের মধ্যে ঔষধ ব্যবহার করতে হবে।
- ঠান্ডা ও শুষ্ক স্থানে এবং শিশুদের নাগালের বাহিরে রাখুন।