Organic Research Food Products Ltd.
Pabna, Bangladesh

Subito Plus

(An effective remedy for primary level of any diseases)

Indication: An effective and fast organic remedy for primary level of any diseases for chicken.

Benefits:

- Protects any diseases of chicken at primary level within 3-4 hours.
- Reduces mortality through rapidly exclusion of pathogens.
- Extremely safe and have no side effect.

Dosage:

- After emerging the first symptoms of any diseases in few chickens, half of powder will be mixed with 100 ml of mineral water and should be provided only one drop orally or half of powder should be mixed with 2-3 litre of mineral water and should be provided for 1000 birds in poultry drinkers. Remaining half of medicine could be used next time after emerging of first symptoms of any diseases.

Precautions:

- This medicine is only for chicks.
- Medicine should be provided after emerging the first symptoms of any diseases in few chickens.
- In case of oral feeding, always use fresh dropper.
- In case of group feeding in drinkers, it should be ensured that all birds have taken the water with medicine.
- Medicine should be used within one month after opening the cap.
- Store in a dry place at a room temperature and keep distant from children.

Organic Research Food Products Ltd.
Pabna, Bangladesh

সুবিতো প্লাস

(প্রাথমিক স্তরের যে কোনো রোগের জন্য একটি কার্যকরী রোগ প্রতিকারক)

কার্যকারিতা: মুরগীর যে কোনো রোগের জন্য একটি কার্যকরী ও দ্রুত জৈব রোগ প্রতিকারক।

উপকারিতা:

- প্রাথমিক অবস্থায় প্রয়োগ করলে ৩-৪ ঘন্টার মধ্যে যে কোনো রোগ প্রতিরোধ করতে পারে।
- দ্রুত জীবানু বর্জনের মাধ্যমে মৃত্যুহার হ্রাস করে।
- অত্যন্ত নিরাপদ ও কোন পার্শ্ব প্রতিক্রিয়া নেই।

ব্যবহার বিধি:

- ✓ খামারে মুরগীর যে কোনো রোগের প্রথম লক্ষণ কয়েকটা মুরগীতে দেখা দেওয়া মাত্রই যে কোনো সময়েই ১০০০ টি মুরগীর জন্য সরবরাহকৃত ঔষধের অর্ধেক পাউডার ১০০ মিলি বোতলজাত পানির সাথে মিশিয়ে শুধুমাত্র এক ফোটা করে প্রতিটি মুরগী/মুরগীর বাচ্চাকে খাওয়াতে হবে অথবা অর্ধেক পাউডার ৮-১০ লিটার বোতলজাত পানির সাথে মিশিয়ে পানির পাত্রে দিয়ে খাওয়াতে হবে। বাকী অর্ধেক ঔষধ পরবর্তীতে কোনো রোগের প্রাথমিক অবস্থায় খাওয়ানোর জন্য রেখে দিতে হবে।

সতর্কতা:

- এই ঔষধ শুধুমাত্র মুরগীর জন্য।
- শুধুমাত্র রোগের প্রাথমিক অবস্থায় প্রয়োগ করতে হবে।
- সরাসরি মুখে খাওয়ানোর ক্ষেত্রে সর্বদা নতুন ড্রপার ব্যবহার করতে হবে।
- পানির পাত্রে ঔষধ খাওয়ানোর ক্ষেত্রে সকল মুরগীর বাচ্চাকে পানি মিশ্রিত ঔষধ খাবার বিষয়টি নিশ্চিত করতে হবে।
- ছিপি খোলার এক মাসের মধ্যে ঔষধ ব্যবহার করতে হবে।
- ঠান্ডা ও শুষ্ক স্থানে এবং শিশুদের নাগালের বাহিরে রাখুন।